



Vision for a Lifetime

CATARACT EYE DROP INSTRUCTIONS LIOL and Regular (Lane)

Please note:

If you have been using eye drops in the **operative eye** (for Glaucoma, etc) keep taking these drops as prescribed up to the day of surgery **BUT do not use** these drops on the day of surgery. Continue normal routine drops in the non- operative **eye**.

DROPS CAN BE USED IN ANY ORDER

SHAKE EYE DROPS BEFORE USING

USE EYE DROPS 5 MINUTES APART

VIGAMOX (Pharmacy may substitute Zymar, Quixin, or IQuix)

- Antibiotic

-Following surgery, use one (1) drop in operative eye three (3) times per day for two (2) weeks.

NEVANAC: (Pharmacy may substitute Xibrom or Acular LS) **or Acuvail**

- Non-steroidal anti-inflammatory.

-Start one (1) day prior to surgery. Use one (1) drop in operative eye three (3) times per day until gone. Use the morning of surgery. **If diabetic; start drops one week prior to surgery.**

- If using **Acuvail**- Use one (1) drop in operative eye two (2) times per day starting the day before surgery. Use the morning of surgery. Continue until gone. **If diabetic; start using one week before surgery.**

**This medication comes in single-use vials.

LOTEMAX or DUREZOL (Diabetics will have Durezol)

- Steroid anti-inflammatory

- Following surgery, use one (1) drop to operative eye three (3) times per day until gone.

BRING ALL BOTTLES OF EYE DROPS WITH YOU TO THE SURGERY CENTER