



Vision for a Lifetime

**CATARACT EYE DROP
INSTRUCTION SHEET
LIOL
Park**

Please note:

If you have been using eye drops in the **operative eye** (for Glaucoma, etc) then you should do the **routine glaucoma drops** in operative eye the day before and morning of surgery **unless told differently** by clinic technician or surgeon.

Continue normal glaucoma drops in the non- operative eye as usual.

**SHAKE EYE DROPS BEFORE USING
USE DROPS 5 MINTUES APART
EYE DROPS CAN BE USED IN ANY ORDER**

VIGAMOX (Pharmacy may substitute Zymar, Quixin, or Iquix)

- Antibiotic

-Start using **after** surgery, use one (1) drop in operative eye three (3) times per day **stop** after two (2) weeks.

NEVANAC (Pharmacy may substitute Xibrom or Acular LS) **or ACUVAIL.**

- Non-steroidal anti-inflammatory.

- If using Nevanac (or substitution) - Use one (1) drop in operative eye three (3) times per day starting the **day before** surgery. Use the morning of surgery. Continue daily **until gone. If diabetic; start using one week before surgery.**

- If using Acuvail- Use one (1) drop in operative eye two (2) times per day starting the day before surgery. Use the morning of surgery. Continue until gone. **If diabetic; start using one week before surgery.** **This medication comes in single-use vials.

Omni-Pred (*Sample will be given to you in the clinic. If you did not receive one call 1-800-846-1877 or 651-275-3000*) If samples no longer available will be ePrescribed Durezol to be used with the same instructions.

- Steroid anti-inflammatory

-Start using **after** surgery, use one (1) drop in operative eye three (3) time per day. Continue until **gone.**

RESTASIS:

- Used to prevent dry eye.

- Start using one week **prior** to surgery.

- Use one (1) drop in operative eye two (2) times per day. Use until supply is gone.

BRING ALL BOTTLES OF EYE DROPS WITH YOU TO THE SURGERY CENTER